Microplastics Are In Our Brains: Here’s What That Means

9 MIN READ

MAR. 17, 2026

BY HEATHER HURLOCK

New research reveals how microplastics may contribute to Alzheimer’s and Parkinson’s.

You already know microplastics are in your **seafood**, your **bottled water**, and your bloodstream. We’ve covered that ground. But a new peer-reviewed study published in ***Molecular and Cellular Biochemistry*** by researchers at the University of Technology Sydney and Auburn University raises a more urgent question: how do microplastics affect brain health?

The short answer is that we don’t fully know yet. The longer answer is worth paying attention to.

Microplastics Linked to Alzheimer’s in New Study

Here’s a quick overview of what the new study found:

ADVERTISEMENT

- **Brain tissue accumulates more microplastics than any other organ**, and levels have

risen 50% between 2016 and 2024.

- **Researchers identified six distinct biological pathways through which microplastics may contribute to Alzheimer’s disease**, and six more linked to Parkinson’s, with several shared between both conditions.
- **No study has yet established direct causation in humans**, and the authors call this out plainly. Most current evidence comes from animal and cell models.
- **Despite the limits of the evidence, the researchers recommend reducing everyday plastic exposure now**, saying the mechanisms are plausible enough, and the changes are low-risk.

The Brain Accumulates More Plastic Than Any Other Organ

A **2025 study in *Nature Medicine*** measured microplastic concentrations in human kidneys, livers, and brains, using autopsy samples collected in 2016 and 2024. Brain tissue held 7 to 30 times more plastic than liver or kidney. Brain concentrations were roughly 50% higher in the 2024 samples than in the 2016 ones, and liver levels rose over the same period. Kidney levels did not.

The same team also analyzed 12 brains from people with a documented dementia diagnosis and found substantially higher concentrations, with deposits clustered in blood vessel walls and immune cells.

ADVERTISEMENT

Here’s where it gets interesting, and where the researchers themselves pump the brakes. They don’t read the dementia finding as evidence that plastic causes dementia. Brain atrophy, a leakier blood-brain barrier, and weakened clearance are all features of dementia itself, and all of them would be expected to let more plastic collect. The arrow may run the other way entirely. The authors also note their detection method is new and not yet standardized across labs.

So what we know is narrower than the headlines suggest: plastic accumulates in brain tissue, and it’s accumulating more than it used to. What it does once it’s there is the open question.

ADVERTISEMENT

The new review maps six specific biological pathways through which microplastics may contribute to Alzheimer’s disease, and six more linked to Parkinson’s. Several are shared between both conditions. These aren’t vague associations. Researchers are now describing how this might be happening at a cellular level.

How Microplastics Affect Brain Health

The blood-brain barrier is the brain’s primary security system, a tightly regulated membrane that controls what enters brain tissue. Microplastics appear to compromise it. Research has shown they do this by disrupting key structural proteins that hold the barrier together. Once that barrier becomes more permeable, inflammatory molecules, immune cells, and additional microplastics can pass through more easily, setting off a cascade of downstream effects.

Associate Professor Kamal Dua of the University of Technology Sydney, one of the study's senior authors, **estimates that adults** now consume roughly 250 grams of microplastics per year, about enough to cover a dinner plate, entering through seafood, salt, processed foods, tea bags, plastic cutting boards, bottled drinks, and even the air we breathe indoors.

From there, the research identifies several mechanisms worth understanding:

ADVERTISEMENT

1. *Neuroinflammation*. The brain's immune cells, called microglia, recognize microplastics as foreign and mount a defensive response. In doing so, they release inflammatory compounds that, over time, can damage the very neurons they're meant to protect. Chronic low-grade neuroinflammation is a well-established feature of both Alzheimer's and Parkinson's disease.
2. *Oxidative stress.* Microplastics appear to increase unstable molecules called reactive oxygen species while simultaneously weakening the body's antioxidant defenses. The brain's dopamine-producing neurons, the ones lost in Parkinson's disease, are particularly vulnerable to this kind of oxidative damage.
3. *Mitochondrial disruption.* Once inside cells, microplastics can travel to the mitochondria and disrupt their function, impairing energy production. In Alzheimer's specifically, reduced energy impairs the transport of tau proteins, contributing to the toxic tangles that are a defining feature of the disease.
4. *Protein misfolding.* In Alzheimer's, abnormal protein clumps (amyloid-beta plaques) and tangled fibers (tau tangles) accumulate and disrupt brain function. In Parkinson's, a

protein called alpha-synuclein misfolds into toxic deposits. A **2023 study published in *Science Advances*** found that nanoplastics accelerate the formation and spread of the toxic protein deposits central to Parkinson's.

5. 5 *The gut-brain connection.* Microplastics disrupt the gut microbiome, increasing intestinal permeability and triggering systemic inflammation. That inflammation can reach the brain via the vagus nerve, the same pathway researchers believe may carry early Parkinson's pathology from the gut upward into the brainstem, consistent with the **Braak hypothesis** of Parkinson's disease progression.
6. 6 *Epigenetic changes.* Perhaps the most overlooked pathway: microplastics appear to alter how genes are expressed in brain cells, downregulating genes that are also known to be suppressed in Alzheimer's disease. This suggests a possible feedback loop, where epigenetic changes increase oxidative stress, which in turn drives further epigenetic disruption.

What the Evidence Can and Can't Tell Us

It's worth being direct about the limits here, because the researchers are. Most of this evidence comes from animal and cell models, not long-term human studies. No study has established a direct causal link between microplastic exposure and Alzheimer's or Parkinson's disease in humans. The authors explicitly call for longitudinal human research, the kind that follows people over years and correlates actual exposure levels with cognitive decline.

There's also a meaningful limitation in the research itself: the majority of studies have used polystyrene microplastics because they're commercially available and easy to work with in lab settings, but polystyrene represents only about 9.7% of global plastic production. More abundant plastics like polyethylene (54.5%) and polypropylene

(16.5%) are far less studied, so the full picture of how different plastic types behave in the brain remains incomplete.

ADVERTISEMENT

What the evidence *does* **support:** microplastics accumulate in the brain, they activate biological pathways we know to be involved in neurodegeneration, and brain concentrations are rising. That's enough to take seriously.

"We need to change our habits and use less plastic. Steer clear of plastic containers and plastic cutting boards, don't use the dryer, choose natural fibers instead of synthetic ones and eat less processed and packaged foods," shares **Dr Keshav Raj Paudel**, a visiting scholar in the UTS Faculty of Engineering, who is studying how inhaled microplastics may affect lung health.

ADVERTISEMENT

Try This: The 5-Point Plastic Exposure Check

You can't eliminate exposure. But some sources contribute far more than others, and the highest-impact reductions are specific and doable. Run through these five areas and note where your exposure is currently highest, that's where to start.

Check your water source. Are you drinking from plastic bottles regularly? Switching to filtered tap water stored in glass or stainless steel is one of the highest-leverage changes available. **Bottled water can contribute up to 90,000 more plastic particles per year** than filtered tap water, with nanoplastics small enough to cross the blood-brain barrier directly. **Here are our favorite water filters.**

Check your food and cookware. Heating food in plastic containers accelerates microplastic leaching. Switch to glass or ceramic for reheating. Plastic cutting boards are also a meaningful source, hardwood or bamboo sheds far fewer particles. **Avoid these foods found to have the most microplastics**.

Check your seafood choices. Filter feeders like mussels and oysters carry the highest microplastic loads. Whole, wild-caught fish with digestive tracts removed carry significantly less. **We've ranked your options here.**

Check your indoor air quality. Synthetic carpets, upholstered furniture, and the tumble dryer all release microfibers into your home's air. A HEPA air purifier reduces what you breathe (We like the **Coway Airmega AP-1512HH Mighty**, which has been shown to reduce pollutants by as much as 99.6% in 30 minutes.) Washing synthetic clothing in a microfiber-catching laundry bag, **Guppyfriend** is a well-tested option, captures particles before they enter the water supply and your air.

Tea bags. **Research has found that a single plastic tea bag** can release billions of microplastic particles into hot water. Loose-leaf tea with a stainless steel infuser is an easy, cost-neutral swap after the initial purchase. (**We like the Reinmoson**.)

*The information provided in this article is for educational and informational purposes only and is not intended as health, medical, or financial advice. Do not use this information to diagnose or treat any health condition. Always consult a qualified healthcare provider regarding any questions you may have about a medical condition or health objectives.**Read our disclaimers**.*

WRITTEN BY:



Heather Hurlock